

## Male Reproduction

As people with CF live longer and healthier lives, many more are becoming parents. Although men with CF can enjoy a normal sex life, most are unable to conceive naturally. With Assisted Reproductive Technologies (ART) many men with CF are able to have their own biological children.

Approximately 98% of men with CF are infertile. This is due to a blocked or absent vas deferens, but sperm are produced normally. The vas deferens is the tube that carries sperm from the testes to the ejaculatory duct. This means that most men with CF will not be able to conceive a child without medical help. However, men with CF still get erections, ejaculate when they orgasm.

### Semen analysis

If you are male with CF, you should consider having a sperm count done whether you are planning a family or not. While there is a 98% chance you don't have any sperm in your semen, there is a 2% chance you do! Semen analysis measures the numbers of sperm, motility (ability to move), size, shape and volume of ejaculate sample. This information can help you to make good sexual and reproductive health decisions now and into the future.

### Assisted Reproductive Technologies

Assisted reproductive technologies (ART) are used if couples are experiencing difficulties conceiving a child naturally. Sperm extraction followed by IVF is the technique which is used to assist men with CF, who are infertile, to conceive a child.

**Sperm Extraction** is a procedure done under local anaesthetic to extract the sperm from either the testicles or the tubes connected to the testicles. Men with CF produce sperm in their testicles, but the blocked or missing vas deferens means that the sperm are unable to get out through the penis. Sperm extraction is carried out so to enable creation of an embryo through IVF.

**Intracytoplasmic sperm injection (ICSI)** is an IVF procedure where a single extracted sperm is directly injected into the woman's egg, assisting with fertilisation and the development of an embryo. This procedure is used in males with CF because of the low ejaculate volume associated with poorly functioning/absent seminal vesicles (structure where semen is produced). The

technique is useful in men with CF because of the problems with semen and sperm.

Usually the female partner will have ovarian stimulation with hormones to increase the number of eggs for collection. The egg collection is usually done under ultrasound procedure. The semen sample is needed on the same day as egg collection.

### Useful Resources

- Assisted reproduction  
[www.rtc.org.au/consumer/index.html](http://www.rtc.org.au/consumer/index.html)
- Family planning <http://srhwa.com.au/>
- Genetic services WA [www.gswa.org.au](http://www.gswa.org.au)
- Female Reproduction Factsheet  
[https://www.cysticfibrosis.org.au/media/wysiwyg/CF-Australia/Fact\\_Sheets/CF\\_Aust\\_Fact\\_Sheet\\_Female\\_Reproduction.pdf](https://www.cysticfibrosis.org.au/media/wysiwyg/CF-Australia/Fact_Sheets/CF_Aust_Fact_Sheet_Female_Reproduction.pdf)
- Pregnancy Factsheet

### Cystic Fibrosis WA

The Niche  
11 Aberdare Rd  
Nedlands WA 6009

Postal Address  
PO Box 959  
Nedlands 6909

**T:** +61 8 6457 7333  
**F:** +61 8 6457 7344  
**E:** [admin@cfwa.org.au](mailto:admin@cfwa.org.au)

Disclaimer: The information contained herein is provided in good faith. However accuracy of any statements is not guaranteed by Cystic Fibrosis Australia. We provide the information on the understanding that persons take responsibility for assessing relevance and accuracy. Individuals are encouraged to discuss their health needs with a health practitioner.

© Copyright Cystic Fibrosis Western Australia 2017