

Pregnancy

One of the most important questions for women with cystic fibrosis (CF) is whether pregnancy will have a detrimental effect on their long-term health. With proper management, careful monitoring, and working closely with both the CF care team and obstetric team, many women with CF are able to carry a child without significantly impacting their long-term health.

There are of course unique demands that pregnancy places on the body, some of which can have serious implications for women with CF. For this reason, women with CF should plan their pregnancies for a time when they are in the best health possible to ensure the most successful outcome.

Am I healthy enough for pregnancy?

Generally, there are four main factors that help determine how well a woman with CF will handle pregnancy. These factors are:

1. Lung function

With a lower FEV1, the risk of complications may be higher. According to experts, lung function is the most important factor that will affect you and your baby's health after delivery.

- *B. cepacia*: Some studies have shown that infection with *Burkholderia cepacia* can lead to more complications during pregnancy. If you have *B. cepacia* and are interested in becoming pregnant, discuss the possible implications for both your and your baby's health this with your CF care team.

2. Nutrition

Ideally, your nutrition status should be good enough to support both you and your baby throughout the pregnancy, including potential morning sickness or changes in appetite. Having a Body Mass Index (BMI) of at least 22 (kg/m²) is a must. Lower BMI is often associated with poor foetal growth and premature delivery.

3. Cystic Fibrosis-Related Diabetes (CFRD)

CFDR increases the risk of problems during pregnancy. Monitoring and controlling blood glucose prior to falling pregnant is very important. Your insulin requirements may change during pregnancy.

4. Cirrhosis (liver disease)

Cirrhosis can be a potential complication in pregnancy for women with CF. Women who have liver disease should discuss the implications of pregnancy with the CF team.

Potential complications during pregnancy

Premature delivery: The most common complication of pregnancy with CF is preterm or premature delivery, which seems to be associated with low pre-pregnancy lung function.

Gestational diabetes: Pregnant women with CF have been found to have a higher risk of gestational diabetes because many already have impaired glucose tolerance. For this reason, your care team should consistently screen for diabetes throughout your pregnancy.

Nutritional deficiencies. Your nutritional demands will change during pregnancy and put you at risk of inadequate weight gain or even weight loss. Most women with CF will need to increase their caloric intake, drink supplements and focus on maintaining and gaining weight while pregnant. It is important to pay close attention to nutritional status and work closely with a CF dietitian throughout pregnancy.

Constipation: Although it is common for all women to experience constipation issues during pregnancy, having CF puts you at a higher risk. To prevent constipation, pregnant women with CF should drink more water, add fibre to their diet and eat more fruits and vegetables. The care team may also recommend the use of stool softeners or laxatives.

Exacerbations: Pregnant women with CF may experience more exacerbations, so it is important to continue your CF respiratory therapies throughout pregnancy to stay as healthy as possible. Your CF care team and obstetrician will want to discuss whether any of your treatments should be stopped or changed during pregnancy, especially if you require antibiotics.

Vitamin A (retinol) toxicity: Because high levels of vitamin A, especially within the first three months, can be associated with developmental issues for your baby, it is important that your care team carefully monitor your vitamin A levels. If you are taking vitamin A supplements, your obstetrician will likely have you stop

taking them during your pregnancy. However, it is important that you do not stop taking your other CF-related vitamins without discussing it with your care team and obstetrician first.

Hypertension: According to some studies, pregnant women with CF have a higher risk of hypertension than those without CF. For this reason, your care team and obstetrician should closely monitor your blood pressure both before and during your pregnancy.

Useful Resources

- CFFood: A Guide for Adults (chapter 10 on pregnancy) www.cfwa.org.au/wp-content/uploads/2017/12/Nutrition-for-Adults.pdf
- Carrier Screening Factsheet www.cfwa.org.au/wp-content/uploads/2018/03/Carrier-Screening-CFCC.pdf
- Female Reproduction Factsheet www.cfwa.org.au/wp-content/uploads/2018/02/CF-Fact-Female-Reproduction.pdf
- Male Reproduction Factsheet www.cfwa.org.au/wp-content/uploads/2018/02/CF-Fact-Male-Reproduction.pdf

Cystic Fibrosis WA

The Niche
11 Aberdare Rd
Nedlands WA 6009

Postal Address
PO Box 959
Nedlands 6909

T: +61 8 6457 7333
F: +61 8 6457 7344
E: admin@cfwa.org.au

Disclaimer: The information contained herein is provided in good faith. However accuracy of any statements is not guaranteed by Cystic Fibrosis WA. We provide the information on the understanding that persons take responsibility for assessing relevance and accuracy. Individuals are encouraged to discuss their health needs with a health practitioner.

© Copyright Cystic Fibrosis Western Australia 2018