

# Cystic Fibrosis Related Diabetes

Cystic Fibrosis Related Diabetes (CFRD) occurs in about 20% of adolescents and 40-60% of adults with cystic fibrosis (CF). CFRD shares features with type 1 and 2 diabetes but is a different condition, with distinct risks and ways to be managed.

One function of the pancreas is to produce insulin; a hormone that helps control the blood sugar, or glucose levels. Glucose is the fuel that the cells need to operate effectively. Insulin is the “key” which allows glucose to enter the cells to be used as energy. However, when the production of insulin is reduced, the cells do not receive the energy they need, and levels of blood glucose rise which can have an adverse effect on health and wellbeing of the individual with CFRD.

CFRD may be triggered by an exacerbation of CF, medication or it could be a gradual onset. With less transport of glucose to the cells, the glucose levels in the blood rise and people will present with:

Lack of energy	A decrease in lung function
Weight loss	Increased hunger
Excessive thirst	Increased urine output

Not everybody with CFRD will have these symptoms, often people are diagnosed during a routine blood test. Some people may experience few symptoms and others have no symptoms at all. For this reason, people with CF are regularly screened for CFRD. The Oral Glucose Tolerance Test (OGTT) is the most sensitive and the best choice for screening for CFRD and it is recommended annually for people with CF who are 10 years and older.

## The importance of screening for CFRD

When symptoms are left untreated, CF symptoms can deteriorate and lead to:

- Increased thickness of mucus which is difficult to clear
- Reduced ability to fight infection
- Loss of muscle mass and loss of weight

Over time, long-term complications, such as damage to small blood vessels, impaired kidney function and vision impairment can occur.

### Pregnancy and CFRD

Women with CF are more prone to developing CFRD due to an increased need to produce extra insulin during pregnancy. When planning a baby, an OGTT is recommended for those who have not been tested within the last 6 months.

### Managing CFRD

Insulin is the treatment of choice for people with CFRD. Insulin helps balance the right amount of glucose in the blood (glycaemic control). When glycaemic control is achieved, people's nutritional status, weight and lung function are improved and their health outcomes optimised. Contrary to people with other types of diabetes, people with CFRD need to continue with the recommended CF nutritional guidelines of a healthy high fat, high energy diet. They should continue to consume food containing carbohydrates throughout the day.

### Low blood sugar (hypoglycemia)

A normal fasting blood glucose level (BGL) is UNDER 5.5 mmol/L. Having a BGL below 4.0mmol/L (hypoglycaemia) can have serious repercussions, because the cells in the body are starved of sugar and not able to function properly. Hypoglycemia could occur when there is a reduction in carbohydrate intake and/or an increase in activity levels, or in response to taking too much insulin. If you are taking medications including insulin for CFRD it is important to eat meals at regular times to help avoid hypoglycaemia.

### Symptoms of hypoglycemia

Symptoms vary from person to person; early signs may include:

Feeling shaky and weak	Sweating
Hunger	Lightheadedness and dizziness
Headache and mood change	Pins and needles around the mouth

### Actions to manage hypoglycemia

Take 15g of fast acting carbohydrates: such as 6-7 small jelly beans, 3 teaspoons of sugar or honey, 150ml of soft drink. Then wait for 15 minutes and retest BGLs:

- If your BGL is still  $\leq$  4.0mmol/L take another 15g of fast acting carbohydrate.
- If your BGL is over 4.0mmol/L, have 15-20g of slow-acting carbohydrate such as a banana, cereal bar or 250ml of milk.

Talk with your diabetes team about how to prevent and manage your hypoglycemia and always carry fast acting carbohydrates with you.

**Important warning**

Please do not drive if your BGLs are below 5.0 mmol/L.

For more information about CFRD, make an appointment to see a Diabetes Educator.

**Useful Resources**

- Cystic Fibrosis Foundation <https://www.cff.org/Life-With-CF/Daily-Life/Cystic-Fibrosis-Related-Diabetes/>
- Women and Children's hospital CFRD booklet <http://www.wch.sa.gov.au/services/az/other/nutrition/documents/CFRDquickguide.pdf>

Special thanks to Sandra Wilberforce, Clinical Nurse – Diabetes Educator, Diabetes Outpatients Centre, Sir Charles Gairdner Hospital for input in to this factsheet.

**Cystic Fibrosis WA**

The Niche  
11 Aberdare Rd  
Nedlands WA 6009

Postal Address  
PO Box 959  
Nedlands 6909

**T:** +61 8 6457 7333  
**F:** +61 8 6457 7344  
**E:** admin@cfwa.org.au

Last revised July 2018

Disclaimer: The information contained herein is provided in good faith. However accuracy of any statements is not guaranteed by Cystic Fibrosis Australia. We provide the information on the understanding that persons take responsibility for assessing relevance and accuracy. Individuals are encouraged to discuss their health needs with a health practitioner.

© Copyright Cystic Fibrosis Western Australia 2017